

Dulcolax



Generic Dulcolax is a stimulant laxative. It acts directly on the bowels, stimulating the bowel muscles to cause a bowel movement. Bisacodyl is used for relieving occasional constipation and irregularity.

Active ingredient: [Bisacodyl](#)

Brand(s):Bisacodyl
Manufacturer: Zydus Cadila Healthcare

Dulcolax 5mg

	Per pill	Per Pack	Order
10 tablet	\$ 0.99	\$ 9.95	Add to cart
30 tablet	\$ 0.66	\$ 19.95 ~~\$29.85~~	Add to cart
60 tablet	\$ 0.63	\$ 37.95 ~~\$59.70~~	Add to cart
90 tablet	\$ 0.61	\$ 54.95 ~~\$89.55~~	Add to cart
120 tablet	\$ 0.58	\$ 69.95 ~~\$119.40~~	Add to cart
180 tablet	\$ 0.50	\$ 89.95 ~~\$179.10~~	Add to cart

Bisacodyl

What is this medicine?

BISACODYL is a stimulant laxative. It acts directly on the bowels, stimulating the bowel muscles to cause a bowel movement. Bisacodyl is used for relieving occasional constipation and irregularity. It may also be used for other conditions as determined by your doctor.

What should I tell my health care provider before I take this medicine?

Tell your health care provider if you have any medical conditions, especially if any of the following apply to you:

- if you are pregnant, planning to become pregnant, or are breast-feeding
- if you are taking any prescription or nonprescription medicine, herbal preparation, or dietary supplement
- if you have allergies to medicines, foods, or other substances
- if you have heart failure, abnormal levels of fluids or electrolytes in your body, nausea, vomiting, or undiagnosed stomach pain

Some MEDICINES MAY INTERACT with bisacodyl. However, no specific interactions with bisacodyl are known at this time.

Ask your health care provider if bisacodyl may interact with other medicines that you take. Check with your health care provider before you start, stop, or change the dose of any medicine.

How should I use this medicine?

Use bisacodyl as directed by your doctor. Check the label on the medicine for exact dosing instructions.

- Take bisacodyl by mouth with or without food.
- Take bisacodyl with a full glass of water (8 oz/240 mL).
- Drinking extra fluids while you are taking bisacodyl is

