

☐

I'm not robot


reCAPTCHA

Continue

Bracon hebetor pdf

Data source: Agricultural Research and Education Organization Agricultural Research, Education and Extension Organization (AREEO) was established by the Act of Parliament on June 30, 1974. The main mandate of the Organization is to develop and manage all research activities of agriculture and natural resources in the Ministry of Agriculture-Jahad. AREEO is determined to facilitate an effective collaboration between Iranian Institutes and World's well recognized research programs in the field. It should be noted that some institutes under the umbrella of AREEO have been established years earlier, such [...] Conference Naderi, G., Seraj, A. A.; Latifian, M.; Takavar, H.; The effects of Bracon hebetor Say as a parasitoid, on Ephestia kuehniella Zeller population fluctuation in date palm stored condition 1100 Naderi, G., Seraj, A. A.; Latifian, M.; Takavar, H. External parasitoid wasp Bracon hebetor Say larvae of some pests stored dates. In this study, change in population growth in four steps but nuts Ephestia kuehniella Zeller affected parasitoid B. hebetor for 6 months in vitro was paid, two populations in this experiment repeated 3 glass box took. First population included 500 pesticides and 50 larvae with parasitoid sex ratio and the second population contains only 500 larvae are pests. Curve changes in population activity was observed two peak density in the first period egg stage, larvae, pupa and insect full 11 weeks, respectively, 6, 8 and 11 weeks during the second 22, 21, 20 and 22 was observed. During 24 week warehousing always pest population growth in four stages assembly conditions (presence of parasitoid population) to less than control treatment (no parasitoid population), respectively, so that the maximum reduction equal to 5.25, 5.36, 4.2 and 4.78 respectively in the four steps were growing pest control suggests that high performance in parasitized by the wasp larvae but nuts and ultimately control it 2012/IR/2012_0.rdf External parasitoid wasp Bracon hebetor Say larvae of some pests stored dates. In this study, change in population growth in four steps but nuts Ephestia kuehniella Zeller affected parasitoid B. hebetor for 6 months in vitro was paid, two populations in this experiment repeated 3 glass box took. First population included 500 pesticides and 50 larvae with parasitoid sex ratio and the second population contains only 500 larvae are pests. Curve changes in population activity was observed two peak density in the first period egg stage, larvae, pupa and insect full 11 weeks, respectively, 6, 8 and 11 weeks during the second 22, 21, 20 and 22 was observed. During 24 week warehousing always pest population growth in four stages assembly conditions (presence of parasitoid population) to less than control treatment (no parasitoid population), respectively, so that the maximum reduction equal to 5.25, 5.36, 4.2 and 4.78 respectively in the four steps were growing pest control suggests that high performance in parasitized by the wasp larvae but nuts and ultimately control it Bibliographic information Publisher: Plant Protection Research Institute All titles: "The effects of Bracon hebetor Say as a parasitoid, on Ephestia kuehniella Zeller population fluctuation in date palm stored condition "Other: Summaries (En, Fa) " " Proceedings of the 19th Iranian Plant Protection Congress: v. 1. pests - v. 2. Plant diseases - v. 3. Weeds, Manzari, Sabbasi, M.Alabadi, FShimi, P., Tehran, Iran: Plant Protection Research Institute, 2010, Iranian Plant Protection Congress, 19th, Tehran, Iran, 2010. - pp. 127 " About AGRIS Contribute Services Acceptable use policy Contact Us About-Shaara HF, Owaysy AA, Ibrahim YV, Basuny NK (2017) A review of impacts of temperature and relative humidity on various activities of honey bees. Insect Conservation and Biodiversity 12(1):1-12. doi:10.1007/s10841-016-9870-8. Subramanyam, D.W. Hightower (eds.), Integrated management of insects in stored products. Marcel Dekker, New York, 1996. Google Scholar Charles AK, George OO, Paul NN, Suresh KR, Ayuka TF (2017) The biology and control of the greater wax moth, Galleria mellonella. Insects, 17, pp 7 Chaudhry SA (1969) Flora of Lylalpur and the adjacent canal -colony districts. West Pakistan. Agric - Uniy Lylalpur, Pakistan: Dabhi R (2011) Comparative biology of Bracon hebetor Say on seven lepidopteran hosts. Karnataka J Agric Sci 24(4):549-550 Google Scholar Duncan BD (1955) Multiple range and multiple F - test. Biometrics 11(1):1-24Article Google Scholar Dutky SR, Thompson JV, Cantwell GE (1964) A technique for the mass propagation of the DD-136 nematode. J Insect Pathol 4:417-422 Google Scholar Ellis JD, Graham JR, Mortensen A (2013) Standard methods for wax moth research. J. Agric. Res. 521-17Article Google Scholar Farag NA, Ismail IA, Elbehery HA, Abdel-Rahman RS, Abdel-Raheem MA (2015) Life table of Bracon hebetor Say. (Hymenoptera: Braconidae) reared on different hosts. Int. J. Chem Technol Res 8(9):123-130 Google Scholar Ghimire MN, Phillips TW (2010) Suitability of different lepidopteran host species for development of Bracon hebetor (Hymenoptera: Braconidae). Environ. Entomol. 39:449-458 Article PubMed Google Scholar Hagstrum DW, Smitthe JB (1977) Host finding ability of Bracon hebetor and its influence upon adult parasite survival and fecundity. Environ Entomol 6:437-439Article Google Scholar Huang CH, Pan XH, Huang DF, Wang BH, Wei GL (2010) Screening of artificial feed formulae for Galleria mellonella L. Guschki Agrivestor Sci (2011) 672-674 Google Scholar Masood AM, Hsin C (2006) Demography of Habrobracon hebetor (Hymenoptera: Braconidae) on Two Pyralid Hosts (Lepidoptera: Pyralidae). Ann Entomol Soc Am 99:102-109 Article PubMed Google Scholar Miller VA, Pappas G, Brower JH, Vail PV, Finn PW (1996) Biological control of stored product insects. In: Subramanyam, D.W. Hightower (eds.), Integrated management of insects in stored products. Marcel Dekker, New York, 1996. Google Scholar Williams JL (1997) Insects: Lepidoptera (moths) In: Morse R., Flottum K., editors. Honey bee pests, predators, and diseases. Al Rout Company, Medina, OH, USA: 121-141 pp.Yones MS, Ma'moun SAM, Farag MA, Abd El-Raouf MN (2019) Hyper spectral application for early diagnosis of American foulbrood disease in the honeybee (Apis mellifera L.) larvae. The Egyptian Journal of Remote Sensing and Space Science 3 Part 2 Month X Mo. of dead larvae/wax combs Infested areas of larvae (inch2)/wax combs No. of infested tunnels/wax combs No. of infested tunnels/wax combs Treated Untreated Treated Untreated September, 2018 1st week 88.0 ± 0.20 13.7 ± 2.52 16.37 ± 1.53 75 2nd week 94.7 ± 1.53 57.35 ± 1.37 87.53 ± 0.58 35 3rd week 93.0 ± 4.02 45.57 ± 35.44 104.83 ± 2.89 85 4th week 90.0 ± 2.0 31.3 ± 0.58 11.0 ± 0.42 95.0 ± 0.95 30 ± 14.07 32.8 ± 12.48 65 ± 5 40.6 ± 1.7 ± 0.72 59 ± 24.47 October 1st week 83.0 ± 0.30 30.0 ± 0.58 15.2 ± 0.20 11.2 week 83.0 ± 3.61 34.17 ± 7.28 89.10 ± 1.13 18 3rd week 93.7 ± 1.58 13rd week 93.7 ± 2.52 42.3 ± 0.58 33.2 ± 1.53 35 4th week 93.5 ± 0.58 3.29 31.27 ± 4.62 18.17 ± 2.08 10 month 88.15 ± 1.899C 34.25 ± 54.4 1.75 ± 1.15 19.0 ± 9.93 ± 0.8 18.5 ± 11.56 November 1st week 88.7 ± 1.53 32.5 ± 4.58 16.0 ± 0.11 2nd week 89.7 ± 0.58 40.14 ± 7.55 53.17 ± 2.89 26 3rd week 87.3 ± 0.30 48.0 ± 0.0 ± 0.21 0.3 ± 0.58 10 4th week 84.7 ± 3.21 29.0 ± 1.15 0.3 ± 0.58 12 November 20.16 ± 0.86 ± 0.58C 14.75 ± 7.54 December 1st week 84.7 ± 2.08 54.4 ± 1.25 4.0 ± 0.0 ± 0.13 2nd week 85.7 ± 3.06 32.37 ± 1.15 27.0 ± 0.0 ± 0.16 3rd week 95.7 ± 1.53 54.4 ± 0.0 ± 0.29 0.0 ± 0.40 4th week 99.3 ± 1.52 82.0 ± 0.0 ± 0.18 November 21.35 ± 0.78 55.5 ± 2.0 8.8 ± 0.29C 19.5 ± 12.30 1.0 ± 1.4D 16.5 ± 3.42 January, 2019 1st week 83.4 ± 4.94 9.9 ± 0.27 2nd week 83.4 ± 4.94 9.9 ± 0.27 3rd week 83.4 ± 4.94 9.9 ± 0.27 4th week 83.4 ± 4.94 9.9 ± 0.27 5th week 83.4 ± 4.94 9.9 ± 0.27 6th week 83.4 ± 4.94 9.9 ± 0.27 7th week 83.4 ± 4.94 9.9 ± 0.27 8th week 83.4 ± 4.94 9.9 ± 0.27 9th week 83.4 ± 4.94 9.9 ± 0.27 10th week 83.4 ± 4.94 9.9 ± 0.27 11th week 83.4 ± 4.94 9.9 ± 0.27 12th week 83.4 ± 4.94 9.9 ± 0.27 13th week 83.4 ± 4.94 9.9 ± 0.27 14th week 83.4 ± 4.94 9.9 ± 0.27 15th week 83.4 ± 4.94 9.9 ± 0.27 16th week 83.4 ± 4.94 9.9 ± 0.27 17th week 83.4 ± 4.94 9.9 ± 0.27 18th week 83.4 ± 4.94 9.9 ± 0.27 19th week 83.4 ± 4.94 9.9 ± 0.27 20th week 83.4 ± 4.94 9.9 ± 0.27 21st week 83.4 ± 4.94 9.9 ± 0.27 22nd week 83.4 ± 4.94 9.9 ± 0.27 23rd week 83.4 ± 4.94 9.9 ± 0.27 24th week 83.4 ± 4.94 9.9 ± 0.27 25th week 83.4 ± 4.94 9.9 ± 0.27 26th week 83.4 ± 4.94 9.9 ± 0.27 27th week 83.4 ± 4.94 9.9 ± 0.27 28th week 83.4 ± 4.94 9.9 ± 0.27 29th week 83.4 ± 4.94 9.9 ± 0.27 30th week 83.4 ± 4.94 9.9 ± 0.27 31st week 83.4 ± 4.94 9.9 ± 0.27 32nd week 83.4 ± 4.94 9.9 ± 0.27 33rd week 83.4 ± 4.94 9.9 ± 0.27 34th week 83.4 ± 4.94 9.9 ± 0.27 35th week 83.4 ± 4.94 9.9 ± 0.27 36th week 83.4 ± 4.94 9.9 ± 0.27 37th week 83.4 ± 4.94 9.9 ± 0.27 38th week 83.4 ± 4.94 9.9 ± 0.27 39th week 83.4 ± 4.94 9.9 ± 0.27 40th week 83.4 ± 4.94 9.9 ± 0.27 41st week 83.4 ± 4.94 9.9 ± 0.27 42nd week 83.4 ± 4.94 9.9 ± 0.27 43rd week 83.4 ± 4.94 9.9 ± 0.27 44th week 83.4 ± 4.94 9.9 ± 0.27 45th week 83.4 ± 4.94 9.9 ± 0.27 46th week 83.4 ± 4.94 9.9 ± 0.27 47th week 83.4 ± 4.94 9.9 ± 0.27 48th week 83.4 ± 4.94 9.9 ± 0.27 49th week 83.4 ± 4.94 9.9 ± 0.27 50th week 83.4 ± 4.94 9.9 ± 0.27 51st week 83.4 ± 4.94 9.9 ± 0.27 52nd week 83.4 ± 4.94

[illegible]