

Aldara



Generic Aldara (Imiquimod) topical is used to treat external warts around the genital and rectal areas called condyloma acuminatum. It is not used on warts inside the vagina, penis, or rectum. Imiquimod is also used to treat a skin condition of the face and scalp called actinic keratosis (AK), which is caused by too much sun exposure. Imiquimod may be used to treat certain types of skin cancer called superficial basal cell carcinoma (sBCC). Imiquimod works on the immune system to help the body fight viruses that cause warts. It does not destroy the viruses directly. It is not known how imiquimod works for actinic keratosis or skin cancer.

Active ingredient: [Imiquimod](#)

Brand(s): Imiquad

Manufacturer: Glenmark

Disease(s): Actinic Keratosis / Skin Cancer

Aldara 5% 0.25g

	Per tube	Per Pack	Order
3 cream	\$ 11.00	\$ 33.00	Add to cart
6 cream	\$ 10.00	\$ 60.00 ~~\$66.00~~	Add to cart
9 cream	\$ 9.50	\$ 85.50 ~~\$99.00~~	Add to cart
12 cream	\$ 9.00	\$ 107.95 ~~\$132.00~~	Add to cart
15 cream	\$ 8.50	\$ 127.45 ~~\$165.00~~	Add to cart
18 cream	\$ 8.00	\$ 143.95 ~~\$198.00~~	Add to cart
24 cream	\$ 7.50	\$ 179.95 ~~\$264.00~~	Add to cart
Free AirMail shipping			
36 cream	\$ 7.00	\$ 251.95 ~~\$396.00~~	Add to cart
Free AirMail shipping			

What is Generic Aldara cream?

Imiquimod topical is used to treat external warts around the genital and rectal areas called condyloma acuminatum. It is not used on warts inside the vagina, penis, or rectum. Imiquimod is also used to treat a skin condition of the face and scalp called actinic keratosis (AK), which is caused by too much sun exposure. Imiquimod may be used to treat certain types of skin cancer called superficial basal cell carcinoma (sBCC).

Imiquimod works on the immune system to help the body fight viruses that cause warts. It does not destroy the viruses directly. It is not known how imiquimod works for actinic keratosis or skin cancer.

Important information

Using too much Aldara, or using it for too long can increase your risk of severe skin reactions. Follow your doctor's instructions.

Do not use Aldara on areas of broken, wounded, or burned skin. Wait until these conditions have healed before using Aldara.

Before using Aldara, tell your doctor if you have a weak immune system, an autoimmune disorder, graft-versus-host disease, or if you have recently received a bone marrow transplant or cord blood transplant.

