

## Lanoxin



Generic Lanoxin is used for treating heart failure and slowing the heart rate in patients with chronic atrial fibrillation, a type of abnormal heart rhythm.

Active ingredient: Digoxin

Brand(s): Lanoxin

Manufacturer: GlaxoSmithKline Plc

Disease(s): Heart Failure / Atrial Fibrillation

### Lanoxin 0.25mg

|            | Per pill | Per Pack                                | Order                       |
|------------|----------|-----------------------------------------|-----------------------------|
| 30 tablet  | \$ 1.10  | <b>\$ 33.00</b>                         | <a href="#">Add to cart</a> |
| 60 tablet  | \$ 0.92  | <b>\$ 55.00</b><br><del>\$66.00</del>   | <a href="#">Add to cart</a> |
| 90 tablet  | \$ 0.83  | <b>\$ 75.00</b><br><del>\$99.00</del>   | <a href="#">Add to cart</a> |
| 120 tablet | \$ 0.75  | <b>\$ 90.00</b><br><del>\$132.00</del>  | <a href="#">Add to cart</a> |
| 180 tablet | \$ 0.67  | <b>\$ 120.00</b><br><del>\$198.00</del> | <a href="#">Add to cart</a> |
| 360 tablet | \$ 0.56  | <b>\$ 200.00</b><br><del>\$396.00</del> | <a href="#">Add to cart</a> |

Free AirMail shipping

## Digoxin tablet

### What is this medicine?

DIGOXIN is used to treat congestive heart failure and heart rhythm problems.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- certain heart rhythm disorders
- heart disease or recent heart attack
- kidney or liver disease
- an unusual or allergic reaction to digoxin, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Take this medicine by mouth with a glass of water. It is best to take this medicine on an empty stomach, at least 1 hour before, or 2 hours after meals. Take your doses at regular intervals. Do not take your medicine more often than directed.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.

NOTE: This medicine is only for you. Do not share this medicine with others.

