

## Lotemax



Generic Lotemax is used for treating swelling, itching, redness, or irritation of the eye caused by bacterial or viral infections, surgery, or certain allergies.

Active ingredient: Loteprednol

**Brand(s):** Loteflam Eye Drops

**Manufacturer:** Cipla Limited

**Disease(s):** Eye Swelling / Eye Redness

### Lotemax 0.5% 5ml

|            | Per solution | Per Pack                              | Order                       |
|------------|--------------|---------------------------------------|-----------------------------|
| 1 solution | \$ 15.00     | <b>\$ 15.00</b>                       | <a href="#">Add to cart</a> |
| 2 solution | \$ 14.00     | <b>\$ 28.00</b><br><del>\$30.00</del> | <a href="#">Add to cart</a> |
| 3 solution | \$ 13.00     | <b>\$ 39.00</b><br><del>\$45.00</del> | <a href="#">Add to cart</a> |
| 4 solution | \$ 12.00     | <b>\$ 48.00</b><br><del>\$60.00</del> | <a href="#">Add to cart</a> |
| 5 solution | \$ 11.00     | <b>\$ 55.00</b><br><del>\$75.00</del> | <a href="#">Add to cart</a> |
| 6 solution | \$ 10.00     | <b>\$ 60.00</b><br><del>\$90.00</del> | <a href="#">Add to cart</a> |

### Loteprednol 0.5% solution

#### What is this medicine?

LOTEPREDNOL is a corticosteroid used in the eye to treat swelling and redness caused by certain eye conditions or surgery.

#### What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions:

- any active infection
- glaucoma
- wear contact lenses
- an unusual or allergic reaction to loteprednol, glycerin, benzalkonium chloride, other corticosteroids, medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

#### How should I use this medicine?

This medicine is only for use in the eye. Do not take by mouth. Shake well before using. Wash your hands before and after use. Tilt your head back slightly, and then pull your lower eyelid down with your index finger to form a pouch. Do not to touch the tip of the dropper to your eye, fingertips, or other surface. Squeeze the prescribed number of drops into the eye pouch. Gently close the eye for a few moments to spread the drops. Do not blink. Keep the eyes closed and apply gentle pressure to the inner corner of the eye with your finger for 1 or 2 minutes to allow the medicine to be absorbed by the eye. Do not use the medicine more often than directed. Do not stop using this medicine earlier than directed by your doctor or health care professional.

