

Afrin



Oxymetazoline is a decongestant that shrinks blood vessels in the nasal passages. Dilated blood vessels can cause nasal congestion (stuffy nose). Oxymetazoline nasal (for the nose) is for temporary relief of nasal congestion (stuffy nose) caused by allergies or the common cold.

Active ingredient: [Oxymetazoline Hydrochloride](#)

Brand(s):Naselin / Zoamet / Oxyspray

Manufacturer: Cipla Ltd. / East India / Zydus Cadila

Disease(s):Allergy

Afrin 0.05% 10ml

	Per spray	Per Pack	Order
3 spray	\$ 9.98	\$ 29.95	Add to cart
6 spray	\$ 6.49	\$ 38.95 ~~\$59.90~~	Add to cart
9 spray	\$ 5.99	\$ 53.95 ~~\$89.85~~	Add to cart

Afrin nasal spray

What is this medicine?

This medication is used for temporary relief of congestion in the nose caused by various conditions including the common cold, sinusitis, hay fever, and allergies. It works by narrowing the blood vessels in the nose area, reducing swelling and congestion.

How should I use this medicine?

Use this medication in the nose as directed. Follow all directions on the product package, or use as directed by your doctor. If you have any questions, ask your doctor or pharmacist. Gently blow your nose before using this drug. Use your finger to close the nostril on the side not receiving the medication. While keeping your head upright, place the spray tip into the open nostril. Spray the medication into the open nostril as you breathe in through your nose. Sniff hard a few times to be sure the medication reaches deep into the nose. Repeat these steps for the other nostril if needed. Avoid spraying the medication into your eyes or onto the middle of the inside of your nose (nasal septum). Rinse the spray tip with hot water or wipe with a clean tissue after each use. Make sure that water does not get inside the container. Replace cap after each use. This medication provides only temporary relief. Do not use more often, use more sprays, or use longer than directed because doing so may increase the risk of side effects. Also, do not use this medication for more than 3 days or it may cause a condition called rebound congestion. Symptoms of rebound congestion include long-term redness and swelling inside the nose and increased runny nose. If this occurs, stop using this medication and consult your doctor or pharmacist. Inform your doctor if your condition worsens or persists after 3 days.

What precautions do I need to take while using this product?

Do not share this medication with others. Use of this container by more than one person may spread infection. Before using this nasal decongestant, tell your doctor or pharmacist if you are allergic to it; or to other sympathomimetics (e.g., pseudoephedrine); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more

